

Macrolepidoptera and Microlepidoptera (Alucitidae and Pyralidae) recorded in Bulgaria, 12 - 24 September 1995 (Lepidoptera)

Stoyan BESHKOV, Barry GOATER

Introduction

During the period September 12-24, 1995 the authors visited several places in SW Bulgaria, East Rhodopi Mts and Black Sea Coast where the lepidopterous fauna was explored. On September 12, a few moths were collected at a 125 W Hg lamp on the balcony of the home of the first of the authors in Sofia, Mladost IV. H.E. In all the other localities given below, moths were collected using a 160 W Hg lamp plugged in, or powered by a Honda EX 350 generator, and by means of a red wine and sugar 1:1 mixture painted on tree trunks. Several species were also found aestivating in tunnels, one in Kresna Gorge and another in Melnik Town, and a few were recorded by day. Nearly all Pyralidae were determined by B. Goater with only a few additions made by S. Beshkov. Alucitidae were identified by S. Beshkov. The only species of Psychidae was determined by P. Hattenschwiler (Uster, Switzerland). All Macrolepidoptera were identified by both authors together, mostly in the field, except for the specimens determined later by examination of genitalia. The identity of all the doubtful species was checked in this way. Only a minimal number specimens were retained, but lists were made of all species seen. The collected material is in the collections of the two authors. Below are included all the data gathered during the expedition. Only a few species of butterflies are listed because public transport (trains and buses) was used for getting about and most of the daylight hours were spent traveling or waiting for transport in a town.

List of the main collecting localities

Kresna - SW Bulgaria, Kresna Gorge, Stara Kresna Railway Station, 200 m alt., 13.IX.1995.

Melnik - SW Bulgaria, Melnik Town, Sandanski District, 400 m alt., 14.IX.1995.
Rozhen - SW Bulgaria, Rozhen Village above Melnik Town, 580 m alt., 15.IX.1995.

Studen Kladenetz - East Rhodopi Mts, Studen Kladenetz Village, Arda Valley, 180 m alt., 18.IX.1995.

Momina Skala - East Rhodopi Mts, Arda Valley, Momina Skala Chalet near Madzharovo Town, 140 m alt., 19.IX.1995.

Meden Buk - East Rhodopi Mts, Zhaltichalsko Dere near Byala Reka River by Meden Buk Village, Ivailovgrad District, 150 m alt., 20.IX.1995.

Byalo Pole (= Belopolyane Village) - East Rhodopi Mts, Ivailovgrad District, 180 m alt., 21.IX.1995.

Arkutino - South Black Sea Coast, "Arkutino" Resting House near Arkutino Swamp and the Mouth of Ropotamo River, Primorsko District, 22.IX.1995.

Balchik (= Balcic) - North Black Sea Coast, Belia Bryag Camping between Balchik and Kavarna towns, 2 km NE from "Tuzlata", 23.IX.1995.

Leucania punctosa (Treitschke, 1828) is new for the Bulgarian fauna. In the list are included 262 species.

In Belia Bryag Camping many species, most of them in considerable numbers, were observed under a *Prunus divariacata* Ldb. tree, feeding on the secretions of the aphid *Pterochloroides persicae* (Chol., 1899), from the family *Lachnidae* (E. Tasheva det.), dropped from the tree. Moths were much more numerous beneath that tree than on the trees which had been anointed with wine-sugar mixture. The species recorded on aphid secretion were: *Udea ferrugalis*, *Nomophila noctuella*, *Camptogramma bilineata*, *Lygephila cracca*, *Hoplodrina ambigua*, *Thalpophila matura*, *Phlogophora meticulosa*, *Atethmia ambusta*, *Discestra trifolii*, *Leucania putrescens*, *Mythimna albipuncta*, *Mythimna vitellina*, *Mythimna l-album*, *Acantholeucania loreyi*, *Noctua pronuba*, *Xestia xanthographa*, *Peridroma saucia*, *Agrotis puta*, *Agrotis ipsilon*, *Agrotis segetum*.

Abbreviations used in the list:

L - specimen/s collected at light

S - specimen/s collected at red wine and sugar 1:1 mixture

D - collected or observed by day

R - collected at rest on lamps in or around buildings

s - single specimen

r - rare species, 2-3 specimens recorded

c - common species, 4-10 specimens recorded

a - abundant, more than 10 specimens recorded

Table 1

List of the species

Names of the taxa	Kresna Melnik Rozhen Studen Momina Meden Byalo Arkutino Balchik Note										
	1	2	3	4	5	6	7	8	9	10	11
HEPIALIDAE											
<i>Triodia amasius dobrogensis</i> (Caradja, 1932) = <i>Korscheltellus amasius ponticus</i> Car. sensu Popescu-Gorj, 1964				L,a	L,r	L,s			L,c		1
<i>Triodia silvina</i> (Linnaeus, 1761)	L,s	L,r	L,s	L,r	L,s	L,s	L,s				2
PSYCHIDAE											
<i>Eochorica balcanica</i> (Rebel, 1919)											3
ALUCITIDAE											
<i>Alucita grammodactyla</i> Zeller, 1841	L,s										4
<i>Alucita cymmatodactyla</i> Zeller, 1852										L,s	5
PYRALIDAE											
<i>Euchromius ocella</i> (Haworth, 1811)										L,s	6
<i>Agriphila tristella</i> ([Denis & Schiffermüller], 1775)											7
<i>Agripigila inquinatella</i> ([Denis & Schiffermüller], 1775)			R,s								8
<i>Agripigila brionella</i> (Zerny, 1914)											9
<i>Agripigila latistria</i> (Haworth, [1811])	D,r,L,s	L,r		L,s							10
<i>Agripigila dalmainella</i> (Hampson, 1900)	D,L,r	L,r		L,c	L,r						11
<i>Pediasia matricella</i> (Treitschke, 1832)	L,c			L,c	L,r						
<i>Catoptiria confusella</i> (Staudinger, 1882)					L,r						
<i>Ancyloloma palpella</i> ([Denis & Schiffermüller], 1775)	L,s	L,r									
<i>Ancylolomia tentaculella</i> (Hübner, 1796)			L,s			L,r					
<i>Eudonia mercurella</i> (Linnaeus, 1758)				L,s							
<i>Evergestia limbata</i> (Linnaeus, 1767)											
<i>Evergestia serratalisi</i> (Staudinger)				L,s	L,c	L,c	L,r			L,s	

	1	2	3	4	5	6	7	8	9	10
<i>Aporodes floralis</i> (Hübner, [1809])		L,s								
<i>Heluta undalis</i> (Fabricius, 1781)						L,s	L,s			
<i>Purausta aurata</i> (Scopoli, 1763)							L,s			
<i>Pyrausta purpuralis</i> (Linnaeus, 1758)		L,s						L,s		
<i>Pyrausta sanguinalis</i> (Linnaeus, 1767)		L,s				L,s				
<i>Pyrausta rectefascialis</i> (Toll, 1936)							L,c			
<i>Ecpyrrhoroe rubiginalis</i> (Hübner, 1796)		L,s		L,s			L,s			
<i>Paracorsia repandalis</i> ([Denis & Schiffermüller], 1775)		L,r		L,s						
<i>Ostrinia nubilalis</i> (Hübner, 1796)		L,s					L,s			
<i>Euctasta splendida</i> (Herrich-Schäffer)		L,s								
<i>Ebulea testacealis</i> (Zeller, 1847)	D,s									
<i>Udea ferrugalis</i> (Hübner, 1796)	L,r	L,r			L,r	L,r	L,r	L,s	L,s;S,r	12
<i>Nomophila noctuella</i> ([Denis & Schiffermüller], 1775)		L,s		L,r	L,s	L,s	L,s	L,s	L,s;S,s	13
<i>Antigastra cataunalis</i> (Duponchel, 1833)						L,r	L,c			
<i>Metasia ophiatis</i> (Treitschke, 1829)	L,r									
<i>Pleuroptiya ruralis</i> (Scopoli, 1763)	L,r	L,r				L,s	L,r			
<i>Palpita unionalis</i> (Hübner, 1796)			L,s		L,s	L,s	L,s			
<i>Palpita incoloralis</i> (Guenée, 1854)					L,s					
<i>Hypsopygia costalis</i> (Fabricius, 1775)										
<i>Actenia brunnealis</i> (Treitschke, 1829)	D,r	L,r								
<i>Actenia honestalis</i> (Treitschke, 1829)										
<i>Orthopygia fulvocitatis</i> (Duponchel, 1834)	L,r							L,s		
<i>Pyralis regalis</i> ([Denis & Schiffermüller], 1775)										
<i>Pyralis farinalis</i> (Linnaeus, 1758)	L,s	R,r							R,r	14
<i>Eudotrica flamealis</i> ([Denis & Schiffermüller], 1775)	L,r	L,s				L,s				
<i>Ocnocera semirubella</i> (Scopoli, 1763)	L,s									
<i>Bradyrrhoa gilveolella</i> (Treitschke)				L,s		L,r	L,s			15
<i>Etiella zinckenella</i> (Treitschke, 1832)		L,r				L,r			L,s	

	1	2	3	4	5	6	7	8	9	10
GEOMETRIDAE										
Hemihelinae										
<i>Hemistola chrysoprasaria</i> (Esper, 1794)		L,s								
<i>Euchrostes indignata</i> (de Villers, 1789)	L,r		L,s							
Sterrhinae										
<i>Cyclophora puppillaria</i> (Hübner, [1799])		L,s		L,s	L,s		L,s			
<i>Timandra griseata</i> W. Petersen, 1902							L,s			
<i>Scopula ochraceata</i> Staudinger, 1901	L,s	L,s		L,r	L,c	L,r	L,r			
<i>Scopula marginipunctata</i> (Goeze, 1781)		L,r		L,r		L,r		L,r	L,s	
<i>Scopula imitaria</i> (Hübner, [1799])	L,s	L,s		L,s					R,s	
<i>Glossotrophia confinaria</i> (Herrich-Schäffer, 1847)		L,s		L,s						23
<i>Idaea filicata</i> (Hübner, [1799])	L,s	L,s								
<i>Idaea dimidiata</i> (Hufnagel, 1767)							L,s			
<i>Idaea camparia</i> (Herrich-Schäffer, [1852])		L,s			L,s			L,s	L,r	24
<i>Idaea aversata</i> (Linnaeus, 1758)								L,s		
<i>Idaea degeneraria</i> (Hübner, [1799])	L,r	L,s		L,s	L,r		L,r			
<i>Rhodometra sacraria</i> (Linnaeus, 1767)					L,s	L,s				
Larentiinae										
<i>Lythria purpuraria</i> (Linnaeus, 1758)	D,s									
<i>Catalysme riguata</i> (Hübner, [1813])		L,s								
<i>Xanthorhoe fluctuata</i> (Linnaeus, 1758)					L,s		L,s			
<i>Epirrhoe galiata</i> ([Denis & Schiffermüller], 1775)									L,s	
<i>Costaconvexa polygrammata</i> (Borkhausen, 1794)										
<i>Campptogramma bilineata</i> (Linnaeus, 1758)		L,s					L,s		R,s,S,s	
<i>Catarhoe permixtaria</i> (Herrich-Schäffer, 1851)	L,r									
<i>Cosmorhoe ocellata</i> (Linnaeus, 1758)									L,s	
<i>Nebula ablutaria</i> (Boisduval, 1840)										
<i>Colostygia</i> sp.							L,s			25
<i>Horysme corticata</i> (Treitschke, 1835)	L,r	L,r							L,r	26

	1	2	3	4	5	6	7	8	9	10
<i>Perizoma bifaciata</i> (Haworth, 1809)	L,s									
<i>Eupithecia linariata</i> ([Denis & Schiffermüller], 1775)								L,s	R,r	
<i>Eupithecia varivirgata</i> Alpheraky, 1878										
<i>Eupithecia contaurvata</i> ([Denis & Schiffermüller], 1775)		L,s					L,s			27
<i>Eupithecia innotata</i> (Hufnagel, 1767)								L,s		28
<i>Eupithecia ochridata</i> Pinker, 1968	L,r						L,s			29
<i>Eupithecia orphnata</i> W. Petersen, 1909					L,s					
<i>Eupithecia ericivata</i> Rambur, 1833							L,s			
<i>Gymnoscelus rufifasciata</i> (Haworth, 1809)				L,s						
<i>Minoa murinata</i> (Scopoli, 1763)	D,r									
<i>Aplocera plagiata</i> (Linnaeus, 1758)									L,s	30
<i>Aplocera derynaria</i> von Mentzer, 1981	L,s	L,s								
Boarmilinae										
<i>Ligdia adustata</i> ([Denis & Schiffermüller], 1775) L,s										
<i>Semiothisa alternaria</i> (Hübner, [1809])			L,s							
<i>Tephrina arnavaecaria</i> ([Denis & Schiffermüller], 1775)										31
<i>Artiora evonymaria</i> ([Denis & Schiffermüller], 1775)							L,r			
<i>Ennomos erosaria</i> ([Denis & Schiffermüller], 1775)										
<i>Ennomos fuscantaria</i> (Haworth, 1809)	L,s					L,s				
<i>Fimera regina</i> Staudinger, 1892						L,s	L,c			
<i>Crocallis tusciaria</i> (Borkhausen, 1793)						L,s	L,r			
<i>Crocallis elinguarina</i> (Linnaeus, 1758)	L,r				L,s					32
<i>Nychiodes dalmatina</i> (Wagner, 1909)	L,r									33
<i>Nychiodes waltheri</i> (Wagner, 1919)							L,c			
<i>Peribatodes rhomboidaria</i> ([Denis & Schiffermüller], 1775)	L,r	L,s	L,s;S,s	L,r	L,r	L,r	L,r	L,s;S,s	L,s;S,s	34
<i>Peribatodes umbraria</i> (Hübner, [1809])	L,r	L,r	L,s	L,r	L,r	L,r	L,r			

	1	2	3	4	5	6	7	8	9	10
<i>Peribatodes correptarius</i> (Zeller, 1847)	L,r			L,r	L,c	L,r	L,r			35
<i>Selidosema plumarium</i> (Denis & Schiffermüller), 1775)	L,r	L,s	L,s			L,r	L,s		L,r	
<i>Gnophos sartatus</i> Treitschke, 1827	L,r			L,s						
<i>Campaca margaritata</i> (Linnaeus, 1767)	L,r			L,s	L,s	L,s	L,r	L,s	R,r;L,r	
<i>Aspitates ochrearia</i> (Rossi, 1794)	D,c;L,r			L,r			L,r	L,r	R,r;L,s	36
<i>Dyscates sicaniaria</i> Zeller, 1852				L,r	L,r					
LASIOCAMPIDAE										
<i>Lasioecampa trifolii</i> (Denis & Schiffermüller), 1775)	L,r	L,r	L,r	L,r	L,c	L,c	L,c		L,s	37
LEMONIIDAE										
<i>Lemonia balcanica</i> (Herrich-Schäffer, 1843)				L,r	L,r		L,r		L,s	
SPHINGIDAE										
<i>Agrilus convolvuli</i> (Linnaeus, 1758)										38
<i>Hyles euphorbiae</i> (Linnaeus, 1758)				L,s			L,s			
<i>Macroglossum stellatarum</i> (Linnaeus, 1758)		D,r		D,c			L,s		L,s	
NOTODONTIDAE										
<i>Pterostoma palpinum</i> (Clerck, 1759)							L,s			
<i>Clostera pygra</i> (Hufnagel, 1766)										
THAUMETOPOEIDAE										
<i>Thaumetopoea pityocampa</i> (Denis & Schiffermüller), 1775)		L,s								
<i>Thaumetopoea solitaria</i> (Freyer, 1838)		L,s				L,r	L,a			
LYMANTRIIDAE										
<i>Lymantria dispar</i> (Linnaeus, 1758)	L,s									
<i>Sphrageidus similis</i> (Fuessly, 1775)					L,s					
ARCTIIDAE										
<i>Eilema sororecula</i> (Hufnagel, 1766)										
<i>Eilema morosina</i> (Herrich-Schäffer, 1848)	L,r	L,r								39
<i>Eilema caniota</i> (Hübner, [1808])	L,s	L,r								40
<i>Eilema pseudocomplana</i> (Daniel, 1939)	L,c				L,c	L,r	L,r			
<i>Phragmatobia fuliginosa</i> (Linnaeus, 1758)			L,s							
<i>Euplagia quadripunctaria</i> (Poda, 1761)	L,s		L,r	L,s						

	1	2	3	4	5	6	7	8	9	10
CTENUCHIDAE										
<i>Dysauxes famula pontica</i> Friese, 1959	L,c	L,c	L,c	L,c	L,c	L,c	L,r		L,s	
NOCTUIDAE										
Hermiiniinae										
<i>Orectis proboscidata</i> (Herrich-Schäffer, [1851])					L,r					
<i>Polyopogon plumigeralis</i> (Hübner, [1825])						S,s				
Rivuliniinae										
<i>Parascotia fuliginaria</i> (Linnaeus, 1761)	L,s		S,s;L,s			L,s				
Hypenodiniinae										
<i>Schrankia costae-strigalis</i> (Stephens, 1834)						L,s		L,r		
Hypeniinae										
<i>Hypena rostralis</i> (Linnaeus, 1758)		L,s;c								41
Catocaliniinae										
<i>Catocala nupta</i> (Linnaeus, 1767)	L,s					S,s				
<i>Catocala elocata</i> (Esper, [1787])		S,s	S,s	S,s		S,s	L,s;S,s			
<i>Catocala puerpera</i> (Giorna, 1791)	S,s									
<i>Grammodes bifasciata</i> (Petagna, 1787)								L,r		
<i>Prodotis stolidia</i> (Fabricius, 1775)				L,s						
<i>Lygephila cracca</i>										
([Denis & Schiffermüller], 1775)										
<i>Apopstes spectrum</i> (Esper, [1787])		S,s							S,s	
<i>Autophila ligaminosa</i> (Eversmann, 1851)		r								42
<i>Tyta luctuosa</i>										
([Denis & Schiffermüller], 1775)		L,s								
Noliniinae										
<i>Meganola strigula</i>										
([Denis & Schiffermüller], 1775)	L,s									
Sarrothripiniinae										
<i>Nycteola siculana</i> (Fuchs, 1899)		L,s							L,s	
<i>Nycteola asiatica</i> (Krulikovskiy, 1904)										
Chloephoriniinae										
<i>Bena bicolorana</i> (Fuessly, 1775)				L,s					L,s	43

	1	2	3	4	5	6	7	8	9	10
<i>Pseudoips prasinanus</i> (Linnaeus, 1758) = <i>P. faganus</i> (Fabricius, 1781)					L,s					44
Acronictinae										
<i>Cryphia algae</i> (Fabricius, 1775)										
<i>Cryphia ochsi</i> Boursin, 1941	S,c;L,r	S,s		S,s						
<i>Cryphia rectilinea</i> (Warren, 1909)	L,s									
<i>Cryphia seladona buryeffi</i> (Draudt, 1931)	L,r			L,r						
<i>Cryphia amasina</i> (Draudt, 1931)				L,s	L,s					45
Acontilinae										
<i>Emmelia trabealis</i> (Scopoli, 1763)										
<i>Acontia lucida</i> (Hufnagel, 1766)		L,s		L,s			L,s			46
<i>Acontia urania</i> Frivaldsky, 1835							L,s			
<i>Eulemma ostrina</i> (Hübner, [1808])		L,s								
<i>Metachrostis velox</i> (Hübner, [1813])		L,s								
Plusiinae										
<i>Euchalcia consona</i> (Fabricius, 1787)							L,s			
<i>Diachrysia chrysiis</i> (Linnaeus, 1758)								L,s		
<i>Macdunnoughia confusa</i> (Stephens, 1850)								L,s		47
<i>Autographa gamma</i> (Linnaeus, 1758)								L,s	L,s;S,s	
<i>Trichoplusia ni</i> (Hübner, [1803])	L,s	L,s		L,s			L,s	L,s		
<i>Abrostola trpartita</i> (Hufnagel, 1766)	L,s									
<i>Abrostola agnorista</i> Dufay, 1956	L,s									48
Cucullinae										
<i>Amphipyra pyramidea</i> (Linnaeus, 1758)	L,s				S,s					49
<i>Amphipyra livida</i> (Denis & Schiffmüller, 1775)										
<i>Amphipyra tragopogonis</i> (Clerck, 1759)		S,s		S,s						
<i>Amphipyra tetra</i> (Fabricius, 1787)		S,r		S,s						
Heliothinae										
<i>Helicoverpa armigera</i> (Hübner, [1808])								L,s	L,s;S,s	
Stirrilinae										
<i>Mycteroplus puniceago</i> (Boisduval, 1840)									R,a	50

	1	2	3	4	5	6	7	8	9	10
Ipimorphinae										
<i>Præstilibia armeniaca</i> Staudinger, 1892	L,c;S,r				L,r		L,c	L,s		51
<i>Platyperigea kadenii</i> (Freyer, 1836)		L,s								
<i>Paradrina flavirena</i> (Guenée, 1852)	L,s	L,r		L,s						52
<i>Paradrina clavipalpis</i> (Scopoli, 1763)					L,s					
<i>Hoplodrina ambigua</i>										
(Denis & Schiffermüller), 1775	L,r	L,r	L,s		L,c	L,r	L,c	L,r	L,s;S,s	53
<i>Spodoptera exigua</i> (Hübner, [1808])	L,s	L,s		L,s		L,r				
<i>Sesamia nonagrioides</i> (Lefebvre, 1827)	L,s									54
<i>Dypterygia scabriuscula</i> (Linnaeus, 1758)	S,s									
<i>Anthraxia eriopoda</i>										
(Herrich-Schäffer, [1851])										55
<i>Mormo maura</i> (Linnaeus, 1758)	L,s	S,s		S,r		S,r				
<i>Polyphaenis subsericata</i>										
(Herrich-Schäffer, [1861])	L,c;S,r			L,s	L,r					
<i>Thalpophila matura</i> (Hufnagel, 1766)	L,r			L,c;S,r	L,r	L,s	L,c;S,r	L,r	R,r;L,s;S,s	
<i>Phlogophora meticulosa</i> (Linnaeus, 1758)		L,s					L,s		S,s	
<i>Mesogona acetosellae</i>										
([Denis & Schiffermüller], 1775)					L,r		L,s			
<i>Atethmia centrigo</i> (Haworth, 1809)							L,s	L,r		
<i>Atethmia ambusta</i>										
([Denis & Schiffermüller], 1775)	L,r									
<i>Agrochola ostheldei</i> Boursin, 1951										
<i>Agrochola gratiosa</i> (Staudinger, 1882)					L,s					56
<i>Episema glaucina</i> (Esper, 1789)										57
<i>Episema tersa</i>										
([Denis & Schiffermüller], 1775)		L,s;S,s	S,r							
<i>Episema ledert</i> Christoph, 1885										
([Denis & Schiffermüller], 1775)		L,s		L,s	L,r		L,r			
<i>Episema leberti</i> Christoph, 1885										
= <i>sareptana</i> Alpheraky, 1897										
<i>Episema korsakovi</i> (Christoph, 1885)										
<i>Aporophila canescens</i> (Duponchel, 1826)										
<i>Meganephria bimaculosa</i> (Linnaeus, 1767)										

	1	2	3	4	5	6	7	8	9	10
<i>Dryobotodes monochroma</i> (Esper, [1790])					L,s					
<i>Dryobotodes carbonis</i> (F. Wagner, 1931)							L,s			
<i>Polymixis serpentina</i> (Treitschke, 1825)					L,r	L,r				
<i>Mesapamea secalis</i> (Linnaeus, 1758)				L,s						58
<i>Luperina rubella rubella</i> (Duponchel, 1835)	L,r							L,r		
<i>Luperina rubella sericea</i> (Caradja, 1932)									L,r	59
<i>Luperina dumertlii</i> (Duponchel, 1826)				L,c	L,r	L,s		L,c		
<i>Gortyna flavago</i> ([Denis & Schiffermüller], 1775)										
Hadeninae										
<i>Discestra trifolii</i> (Hufnagel, 1766)		L,r		L,s		L,s	L,r	L,s	S,s	
<i>Lacanobia oleracea</i> (Linnaeus, 1758)			S,s				L,s			60
<i>Leucania herrichi</i> Herrich-Schäffer, 1849				L,r						
<i>Leucania putrescens</i> (Hübner, [1824])	L,s;S,r	L,s	L,s;S,r				S,s		S,r	61
<i>Leucania punctosa</i> (Treitschke, 1828)										
<i>Mythimna ferrago</i> (Fabricius, 1787)										
<i>Mythimna albipuncta</i>										
([Denis & Schiffermüller], 1775)	L,r									
<i>Mythimna vitellina</i> (Hübner, [1808])				L,r;S,s	L,r		L,r	L,s	L,s;S,s	
<i>Mythimna congrua</i> (Hübner, [1817])				L,r;S,s	L,s	L,s	L,r	L,s	R,s;L,s;S,s	
<i>Mythimna l-album</i> (Linnaeus, 1767)									L,s	
<i>Mythimna scirpi</i> (Duponchel, 1836)	L,s	L,r							R,s;L,s;S,s	
<i>Acantholeucania loreyi</i> (Duponchel, 1827)										
<i>Tholera decimatis</i> (Poda, 1761)					L,s	L,s			L,s;S,s	
Noctulinae										
<i>Ochroleura flammatra</i>										
([Denis & Schiffermüller], 1775)	L,s									
<i>Noctua pronuba</i> Linnaeus, 1758	S,s;L,r			S,s	L,s	L,r	L,r		S,s	
<i>Latanoctua orbona</i> (Hufnagel, 1766)									S,s	
<i>Paranoctua commes</i> (Hübner, [1813])	L,r		L,s	L,s	L,s;S,s	L,s;S,s	L,s			
<i>Paranoctua interposita</i> (Hübner, [1870])									L,s	62
<i>Lampra fimbriata</i> (Schreber, 1759)				L,s	L,r					63

	1	2	3	4	5	6	7	8	9	10
<i>Lampra tirrenica</i> (Biebinger, Speidel & Hanigk, 1983)	L,s	L,s								64
<i>Euschestis tertia</i> (von Mentzer, Moberg & Fibiger, 1991)			S,s							65
<i>Internoctua interjecta</i> (Hübner, [1803])	L,s;S,s						L,s	L,r		
<i>Divana haywardi</i> (Tams, 1926)										
<i>Epilecta linogrisea</i> ([Denis & Schiffermüller], 1775)				L,s	L,r					
<i>Chersotis rectangularis</i> ([Denis & Schiffermüller], 1775)		L,s			L,r					66
<i>Eugnorisma pontica</i> (Staudinger, 1892)		L,r;S,c	S,a							
<i>Xestia c-nigrum</i> (Linnaeus, 1758)	L,s							L,s		
<i>Xestia xanthographa</i> ([Denis & Schiffermüller], 1775)	L,s;S,r	S,c;L,c	S,s;L,s	L,s	L,r		L,s;S,s	L,c	L,s;S,s	67
<i>Xestia cohaesa</i> (Herrich-Schäffer, [1849])		L,c			L,a		L,c		S,r	68
<i>Peridroma saucia</i> (Hübner, [1808])									L,s	
<i>Euxoa cos</i> (Hübner, 1824)										
<i>Euxoa distinguenda distincta</i> Staudinger, 1892										
<i>Euxoa temera</i> (Hübner, [1808])									L,r	
<i>Euxoa segnifis</i> (Duponchel, 1836)									L,s	
<i>Euxoa obelisca</i> ([Denis & Schiffermüller], 1775)									L,s	
<i>Agrotis obesa scita</i> Alphéraky, 1889 = <i>A. obesa nivea</i> Caradja, 1932 (L.T.: Balciç)			L,s		L,s					
<i>Agrotis crassa</i> (Hübner, [1803])	S,r;L,r	L,r;S,r	L,s;S,s	L,r		L,s				
<i>Agrotis puta</i> (Hübner, [1803])	L,r;S,r		L,s						L,r;S,s	
<i>Agrotis ipsilon</i> (Hufnagel, 1766)	L,s					S,s			L,s;S,s	
<i>Agrotis trux</i> (Hübner, [1824])	L,c;S,r	L,r;S,r		L,s;S,s	L,s					
<i>Agrotis segetum</i> ([Denis & Schiffermüller], 1775)	S,r	S,s					S,s		S,s	
<i>Agrotis vestigialis</i> (Hufnagel, 1766)										
<i>Agrotis exclamatoris</i> (Linnaeus, 1758)									L,a	69

Notes

1. L.T.: - Balcic (= Balchik), 08. and 20. November, 1931 and Carmen Silva (Romania) - CARADJA (1932: 41) - *Hepialus dobrogensis* ssp. (?) nov. and p. 45: *Hepialus amasinus* ssp. *dobrogensis*. In some articles GANEV (1984: 416) and some others the year of the description of the species is wrongly given as 1934 (Dt. Ent. Z. Iris, Bd. 48: 185-191). Genitalia of many specimens from Balchik and East Rhodopi Mts checked.

2. In POPESCU-GORJ (1964) in the text (p. 19) this species is given as *Korscheltellus amasinus dobrogensis* Car., but in the illustration (Fig. 1.) as *Korscheltellus amasinus ponticus* Car. We consider *Korscheltellus amasinus ponticus* sensu POPESCU-GORJ (1964) as a syn. n., nomen nudum of *Triodia amasinus dobrogensis* (Caradja, 1932).

3. The genitalia of all *Triodia* specimens in bad condition were checked.

4. More than 20 ♂ specimens at light, det. P. Hattenschwiler (Uster, CH).

5. One ♂, genitalia checked.

6. Genitalia checked.

7. Genitalia checked.

8. Rozhenski Manastir, 15.IX.1995, one inside building.

9. Genitalia checked.

10. Genitalia checked.

11. Genitalia checked from all localities.

12. Sofia Town, "Mladost IV" H.E., 12.IX.1995.

13. Sofia Town, "Mladost IV" H.E., 12.IX.1995.

14. Several specimens observed on a ceiling in a market in Melnik Town.

15. Genitalia of all specimens checked.

16. Genitalia checked.

17. East Rhodopi Mts, Haskovo Town, 18.IX.1995, single specimen.

18. Single specimen observed near Novo Delchevo Village in SW Bulgaria, Melnik District, 14.IX.1995.

19. Wings of specimens eaten by bats in the tunnel under Stara Kresna Railway Station; 11 specimens observed aestivating in the tunnel in the museum "Kordopulovata Kashta" House in Melnik Town, 15.IX.1995.

20. Wings of specimens eaten by bats in the tunnel under Stara Kresna Railway Station.

21. Central Bulgaria, Stara Zagora Town, 22.IX.1995, several specimens observed.

22. Genitalia checked.

23. Genitalia checked.

24. Genitalia checked by both authors.

25. One ♂ and one ♀ (undetermined), Gen. preps 21-22./30.I.1996, Beshkov.

26. Rozhenski Manastir, 15.IX.1995, three specimens inside building.




Fig. 1 - 17.

First column: 1. *Triodia amasinus dobrogensis*, ♀ - Balchik, 23.IX.1995. 2. *Triodia amasinus dobrogensis*, ♂ - Balchik, 23.IX.1995. 3. *Eochorica balcanica*, ♂ - Melnik, 14.IX.1995. 4. *Eupithecia variostrigata*, ♀ - Balchik, 23.IX.1995. 5. *Aplocera dervenaria*, ♀ - Melnik, 14.IX.1995. 6. *Dyscia sicanaria*, ♂ - Byalo Pole, 21.IX.1995.

Second column: 7. *Eilema pseudocomplana*, ♀ - Byalo Pole, 21.IX.1995. 8. *Eilema caniola*, ♂ - Byalo Pole, 21.IX.1995. 9. *Mycteroplus puniceago*, ♂ - Balchik, 23.IX.1995. 10. *Praestilbia armeniaca*, ♂ - Byalo Pole, 21.IX.1995. 11. *Episema lederi*, ♂ - Byalo Pole, 21.IX.1995. 12. *Luperina rubella sericea*, ♀ - Balchik, 23.IX.1995.

Third column: 13. *Eugnorisma pontica*, ♂ - Melnik, 14.IX.1995. 14. *Eugnorisma pontica*, ♂ - Melnik, 14.IX.1995. 15.

Xestia cohaesa, ♀ - Momina Skala Chalet near Madzharovo, 19.IX.1995. 16. *Agrotis obesa scytha*, ♂ - Balchik, 23.IX.1995. 17. *Agrotis vestigialis*, ♂ - Arkutino, 22.IX.1995.

27. One ♀, genitalia checked.

28. Collected also in Sofia Town, "Mladost IV" H.E., 12.IX.1995. Genitalia of all specimens checked.

29. One ♂ specimen, genitalia checked.

30. Genitalia checked.

31. SW Bulgaria, Gradeshnitsa Village, Kresna District, 14.IX.1995, one on window of inn.

32. Genitalia not checked.

33. Genitalia not checked.

34. East Rhodopi Mts, Borislavtsi Village, 20.IX.1995, single specimen resting on a wall.

35. SW Bulgaria, Gradeshnitsa Village, Kresna District, 14.IX.1995, one on window of inn.

36. Ivailovgrad Dam, 20.IX.1995, three specimens resting on a wall by police checkpoint.

37. East Rhodopi Mts, Borislavtsi Village, 20.IX.1995, single specimen resting on a wall.

38. East Rhodopi Mts, Madzharovo Town, 19.IX.1995, one resting on a post.

39. Genitalia of many specimens checked.

40. Genitalia of many specimens checked.

41. Approximately 10 specimens aestivating in the tunnel under Stara Kresna Railway Station; four specimens observed aestivating in the tunnel in the museum "Kordopulovata Kashta" House in Melnik Town, 14.IX.1995.

42. Three specimens collected aestivating in the tunnel in the museum "Kordopulovata Kashta" House in Melnik Town, 14.IX.1995.

43. For the nomenclature see MIKKOLA & HONEY (1993).

44. For the nomenclature see MIKKOLA & HONEY (1993).

45. Both specimens worn.

46. Sofia Town, "Mladost IV" H.E., 12.IX.1995.

47. One ♂ specimen, Genitalia checked.

48. For the nomenclature see MIKKOLA & HONEY (1993).

49. Genitalia checked.

50. 29 specimens collected at rest inside buildings to which they had been attracted by previous nights' illumination. The day before the collecting the temperature had decreased by about 15°C.

51. Rozhenski Manastir, 15.IX.1995, several males inside building.

52. Sofia Town, "Mladost IV" H.E., 12.IX.1995.

53. Sofia Town, "Mladost IV" H.E., 12.IX.1995.

54. Rozhenski Manastir, 15.IX.1995, one inside building.

55. One worn specimen at sugar.

56. East Rhodopi Mts, Borislavtsi Village, 20.IX.1995, single specimen resting on a wall.

57. In Bulgaria known from Kresna Gorge (GANEV & HACKER, 1984; BESHKOV, 1992), East Rhodopi Mts: Studen Kladenetz Village and Momina Skala Chalet near Madzharovo Town (BESHKOV, 1995).

58. Genitalia of all *Mesapamea* specimens checked.

59. L.T.: - Balcic (Balchik), 06.November, 30. December 1930 CARADJA (1932) - *Palluperina (Luperina) rubella* ssp. nov. *sericea*.

60. Two ♀ specimens in a bad condition were collected. Till now known in Bulgaria only from East Rhodopi Mts, Momina Skala Chalet near Madzharovo (BESHKOV, 1995).

61. Single male specimen (Fig. 18) at sugar, a new species for the Bulgarian fauna.

62. One worn specimen, genitalia checked.



Fig. 18. *Leucania punctosa*, ♂ - Byalo Pole, 21.IX.1995.

63. Genitalia checked.

64. Genitalia checked.

65. Sofia Town, "Mladost IV" H.E., 12.IX.1995.

66. All specimens worn.

67. Genitalia of many specimens checked.

68. Genitalia checked.

69. Sofia Town, "Mladost IV" H.E., 12.IX.1995.

Acknowledgments

We would like to express our sincere gratitude to Mr P. Hattenschwiler (Uster, Switzerland) for determination of Psychidae species and to Dr. E. Tasheva (Sofia) for the determination of the aphid species from Belia Bryag.

References

- CARADJA A. 1932. Beiträge zur Lepidopteren-Fauna Grossrumäniens für das Jahr 1931. I. Südliche Dobrogea. - Bull. Sect. Scient. Acad. Roum., 15 (1-2): 35-45.
- CARADJA A. 1932. Neuer Beiträge zur Kenntnis der Lepidopteren-Fauna Rumäniens. - Dt. Ent. Z. Iris, 48: 185-191.
- BESHKOV S. 1995. A contribution to the knowledge of the Bulgarian Lepidoptera fauna (Lepidoptera: Macrolepidoptera). - Phegea, 23 (4): 201-218.
- BESHKOV S. 1992. Faunistic Advances on Bulgarian Lepidoptera. - Boll. Ass. Romana Entomol., 46: 37-56.
- FIBIGER M., H. HACKER. 1990. Systematic List of the Noctuidae of Europe. - Esperiana, 2: 1-109.
- FIBIGER M., H. HACKER. 1992. Systematic List of the Noctuidae of Europe. Corrigenda et Addenda I. - Esperiana, 3: 507-511.
- GANEV J. 1984. Catalogue of the Bulgarian Bombyces and Sphingines (Lepidoptera: Notodontidae, Dilobidae, Thaumetopoeidae, Ctenuchidae, Saturniidae, Endromidae, Lasiocampidae, Sphingidae, Hepialidae, Cossidae, Thyrididae, Limacodidae, Drepanidae, Thyatiridae, Lymantriidae, Arctiidae, Nolidae). - Entomofauna, 5 (33, 1-2): 391-467.
- GANEV J., H. HACKER. 1984. *Agrochola osthelderi* Boursin, 1951 eine neue Art für Europa (Lepidoptera, Noctuidae, Cuculliinae). - Articulata, 2 (5): 101-105.
- MIKKOLA K., M. HONEY. 1993. The Noctuoidea (Lepidoptera) described by Linnaeus. - Zool. Journ. Linn. Soc., 108: 103-169.
- POPESCU-GORJ A. 1964. Catalogue de la collection de Lépidoptères "Prof. A. Ostrogovich" du Muséum d'histoire naturelle "Grigore Antipa", Bucarest. Bucarest, Mus. Hist. Nat. "Gr. Antipa". 281 p., pl. XVIII.

Received on 3.11.1998

Addresses of the authors:

Stoyan Beshkov
National Museum of Natural History
1, Tsar Osvoboditel Blvd
1000 Sofia, Bulgaria

Barry Goater
"The Ridge", 27 Hiltingbury Road
Chandlers's Ford, Eastleigh,
Hants, SO53 5SR, England

Macrolepidoptera u Microlepidoptera (Alucitidae u Pyralidae), установени в България в периода 12 - 24 септември 1995 (Lepidoptera)

Стоян БЕШКОВ, Бари ГОТЪР

(Р е з ю м е)

Представени са резултатите от колекционната експедиция на авторите през споменатия период в Югозападна България, Източните Родопи, Черноморското крайбрежие и някои други части на страната. Във всичките споменати в работата находища нощните пеперуги са ловени на светлинна ловилка и на хранителна примамка от червено вино и захар. За всеки от съобщените видове е указан начинът и времето на улавянето му и приблизителният период на наблюдаваните екземпляри. В списъка са включени всичките 262 вида, установени през периода. *Leucania punctosa* (Treitschke, 1828) е нов вид за пеперудната фауна в България. Голяма част от видовете са нови за изследваните райони, друга част са с ограничен брой находища в страната. Приложени са снимки на новия и на останалите по-интересни видове и някои допълнителни пояснения.